



PCT/AU2004/001416

Patent Office
Canberra

I, LEANNE MYNOTT, MANAGER EXAMINATION SUPPORT AND SALES hereby certify that annexed is a true copy of the Provisional specification in connection with Application No. 2003905726 for a patent by INTER-K PTY LIMITED as filed on 17 October 2003.



WITNESS my hand this
Twenty-ninth day of October 2004

A handwritten signature in black ink, appearing to be 'dA' or similar, written over a horizontal line.

LEANNE MYNOTT
MANAGER EXAMINATION SUPPORT
AND SALES

BEST AVAILABLE COPY

AUSTRALIA
Patents Act 1990 (Cth)

PROVISIONAL SPECIFICATION

INTER-K Pty Limited

Invention Title

Methods and Agents for the Treatment of Cancer

The invention is described in the following statement:

Blake Dawson Waldron Patent Services
Level 41, Grosvenor Place
225 George Street
SYDNEY NSW 2000
Telephone: + 61 2 9258 6000
Fax: + 61 2 9258 6999

Ref: DGC DAA 1346 5727

Methods And Agents For The Treatment Of Cancer

FIELD OF THE INVENTION

The present invention relates to methods for modulating cell activity mediated by mitogen activated protein kinases (MAP's) and more particularly, inhibition of the growth and/or the proliferation of cancer cells. There are also provided agents for use in the methods of the invention.

BACKGROUND OF THE INVENTION

The spread of cancer cells involves tumour cell migration through the extracellular matrix scaffold, invasion of basement membranes, arrest of circulating tumour cells, and tumour cell extravasation and proliferation at metastatic sites. Detachment of cells from the primary tumour mass and modification of the peri-cellular environment aid penetration of tumour cells into blood and lymphatic vessels. It is the invasive and metastatic potential of tumour cells that ultimately dictates the fate of most patients suffering from malignant diseases. Hence, tumourigenesis can be viewed as a tissue remodelling process that reflects the ability of cancer cells to proliferate and digest surrounding matrix barriers. These events are thought to be regulated, at least in part, by cell adhesion molecules and matrix-degrading enzymes.

Cell adhesion receptors on the surface of colon cancer cells are involved in complex cell signalling which may regulate cell proliferation, migration, invasion and metastasis and several families of adhesion molecules have now been identified including integrins, cadherins, the immunoglobulin superfamily, hyaluronate receptors, and mucins. In general, these cell surface molecules mediate both cell-cell and cell-matrix binding, the latter involving attachment of tumour cells to extracellular scaffolding molecules such as collagen, fibronectin and laminin.

Of all the families of cell adhesion molecules, the best-characterised is the family known as integrins. Integrins are involved in several fundamental processes including leucocyte recruitment, immune activation, thrombosis, wound healing, embryogenesis, virus internalisation and tumourigenesis. Integrins are transmembrane glycoproteins consisting of an alpha (α) and beta (β) chain in close association that provide a structural and functional bridge between extracellular matrix molecules and cytoskeletal

components with the cell. The integrin family comprises 17 different α and 8 β subunits, and the $\alpha\beta$ combinations are subsumed under 3 subfamilies.

Excluding the leucocyte integrin subfamily that is designated by the $\beta 2$ nomenclature, the remaining integrins are arranged into two major subgroups, designated $\beta 1$ and αv based on sharing common chains.

In the $\beta 1$ subfamily, the $\beta 1$ chain combines with any one of nine α chain members ($\alpha 1-9$), and the α chain which associates with $\beta 1$ determines the matrix-binding specificity of that receptor. For example, $\alpha 2\beta 1$ binds collagen and laminin, $\alpha 3\beta 1$ binds collagen, laminin and fibronectin, and $\alpha 5\beta 1$ binds fibronectin. In the αv subfamily of receptors, the abundant and promiscuous αv chain combines with any one of five β chains, and a distinguishing feature of αv integrins is that they all recognise and bind with high affinity to arginine-glycine-aspartate (RGD) sequences present in the matrix molecules to which they adhere.

The current picture of integrins is that the N-terminal domains of α and β subunits combine to form a ligand-binding head. This head, containing the cation binding domains, is connected by two stalks representing both subunits, to the membrane-spanning segments and thus to the two cytoplasmic domains. The β subunits all show considerable similarity at the amino acid level. All have a molecular mass between 90 and 110 kDa, with the exception of $\beta 4$ which is larger at 210 kDa. Similarly, they all contain 56 conserved cysteine residues, except for $\beta 4$ which has 48. These cysteines are arranged in four repeating patterns which are thought to be linked internally by disulphide bonds. The α -subunits have a molecular mass ranging from 150 - 200 kDa. They exhibit a lower degree of similarity than the β chains, although all contain seven repeating amino acid sequences interspaced with non-repeating domains.

The β subunit cytoplasmic domain is required for linking integrins to the cytoskeleton. In many cases, this linkage is reflected in localisation to focal contacts, which is believed to lead to the assembly of signalling complexes that include α -actinin, talin, and focal adhesion kinase (FAK). At least three different regions that are required for focal contact localisation of $\beta 1$ integrins have been delineated (Reszka et al, 1992). These regions contain conserved sequences that are also found in the cytoplasmic domains of the $\beta 2$, $\beta 3$, $\beta 5$, $\beta 6$ and $\beta 7$ integrin subunits. The functional differences between

these cytoplasmic domains with regard to their signalling capacity have not yet been established.

The integrin $\beta 6$ subunit was first identified in cultured epithelial cells as part of the $\alpha v \beta 6$ heterodimer, and the $\alpha v \beta 6$ complex was shown to bind fibronectin in an arginine-glycine-aspartate (RGD)-dependent manner in human pancreatic carcinoma cells (Sheppard et al, 1990). The $\beta 6$ subunit is composed of 788 amino acids and shares 34 - 51% sequence homology with other integrin subunits $\beta 1 - \beta 5$. The $\beta 6$ subunit also contains 9 potential glycosylation sites on the extracellular domain (Sheppard et al, 1990). The cytoplasmic domain differs from other subunits in that it is composed of a 41 amino acid region that is highly conserved among integrin subunits, and a unique 11 amino acid carboxy-terminal extension. The 11 amino acid extension has been shown not to be necessary for localisation of $\beta 6$ to focal contacts. In fact, its removal appears to increase receptor localisation. However, removal of any of the three conserved regions identified as important for the localisation of $\beta 1$ integrins to focal contacts (Reszka et al, 1992) has been shown to eliminate recruitment of $\beta 6$ to focal contacts (Cone et al, 1994).

The integrin $\alpha v \beta 6$ has previously been shown to enhance growth of colon cancer cells *in vitro* and *in vivo*, and this growth-enhancing effect is due, at least in part, to $\alpha v \beta 6$ mediated gelatinase B secretion (Agrez et al, 1999). What has made this epithelial-restricted integrin of particular interest in cancer is that it is either not expressed or expressed at very low levels on normal epithelial cells, but is highly expressed during wound healing and tumourigenesis, particularly at the invading edge of tumour cell islands (Breuss et al, 1995; Agrez et al, 1996).

Integrins can signal through the cell membrane in either direction. The extracellular binding activity of integrins can be regulated from the cell interior as, for example, by phosphorylation of integrin cytoplasmic domains (inside-out signalling), while the binding of the extracellular matrix (ECM) elicits signals that are transmitted into the cell (outside-in signalling) (Giancotti and Ruoslahti, 1999). Outside-in signalling can be roughly divided into two descriptive categories. The first is 'direct signalling' in which ligation and clustering of integrins is the only extracellular stimulus. Thus, adhesion to ECM proteins can activate cytoplasmic tyrosine kinases (eg. focal adhesion kinase FAK) and serine/threonine kinases (such as those in the mitogen-activated protein kinase (MAPK) cascade) and stimulate lipid metabolism (eg. phosphatidylinositol-4, 5-

biphosphate (P_iP_2) synthesis. The second category of integrin signalling is 'collaborative signalling', in which integrin-mediated cell adhesion modulates signalling events initiated through other types of receptors, particularly receptor tyrosine kinases that are activated by polypeptide growth factors (Howe et al, 1998). In all cases, however, integrin-mediated adhesion seems to be required for efficient transduction of signals into the cytosol or nucleus.

MAP kinases behave as a convergence point for diverse receptor-initiated signalling events at the plasma membrane. The core unit of MAP kinase pathways is a three-member protein kinase cascade in which MAP kinases are phosphorylated by MAP kinase kinases (MEKs) which are in turn phosphorylated by MAP kinase kinase kinases (e.g. Raf-1) (Garrington and Johnson, 1999). Amongst the 12 member proteins of the MAP kinase family are the extracellular signal-regulated kinases (ERKs) (Boulton et al, 1991) activated by phosphorylation of tyrosine and threonine residues which is the type of activation common to all known MAP kinase isoforms. ERK 1/2 (44kD and 42kD MAPks, respectively) share 90% amino acid identity and are ubiquitous components of signal transduction pathways (Boulton et al, 1991). These serine/threonine kinases phosphorylate and modulate the function of many proteins with regulatory functions including other protein kinases (such as $p90^{msk}$) cytoskeletal proteins (such as microtubule-associated phospholipase A_2), upstream regulators (such as the epidermal growth factor receptor and Ras exchange factor) and transcription factors (such as c-myc and Elk-1). ERKs play a major role in growth-promoting events, especially when the concentration of growth factors available to a cell is limited (Giancotti and Ruoslahti, 1999).

Recently, MAP kinases have been found to associate directly with the cytoplasmic domain of integrins, and the binding domains of $\beta 3$, $\beta 5$ and $\beta 6$ for ERK2 have been characterised (see International Patent Application No. WO 01/000677 and International Patent Application No. WO 02/051993). Those patent applications also showed that the cellular activity of cancer cells expressing $\beta 6$ can be modulated by inhibiting binding of the MAP kinase with the integrin by treating the cells with peptides comprising the binding domain for the MAP kinase linked to the carrier peptide penetratin.

The distribution of $\beta 6$ integrin subunit within various tissues has been assessed by both *in situ* hybridisation and immunostaining and reported in the art. For instance, $\beta 6$ mRNA in adult primate tissues was detected only in epithelial cells and at very low or

undetectable levels in most normal tissues (Breuss et al, 1993). High-level expression of $\beta 6$ has been observed in secretory endometrial glands while low-level expression was detected in the ductal epithelia of salivary gland, mammary gland and epididymis, in gall and urinary bladder, and in the digestive tract.

5 Immunostaining data have also shown that $\beta 6$ expression is restricted to epithelia and is up-regulated in parallel with morphogenetic events, tumourigenesis and epithelial repair (Breuss et al, 1993; 1995). During development of the kidney, lung and skin, $\beta 6$ is expressed by specific types of epithelial cells, whereas it is mostly undetectable in normal adult kidney, lung and skin. In contrast, high level expression of $\beta 6$ has been observed in
10 several types of carcinoma. For example, $\beta 6$ is almost invariably neo-expressed in squamous cell carcinomas derived from the oral mucosa, and often focally localised at the infiltrating edges of tumour cell islands (Breuss et al, 1995). Moreover, expression of the $\beta 6$ subunit has been observed in renal cell carcinoma and testicular tumour cell lines (Takiuchi et al, 1994) and 50% of lung cancers have been shown to express this subunit
15 (Smythe et al, 1995).

Recent studies have also shown that $\alpha v \beta 6$ is a major fibronectin-binding receptor in colorectal cancer (Agrez et al, 1996). In addition, normal colonic epithelium from cancer patients does not express $\alpha v \beta 6$ in immunostaining studies, and as with squamous cell carcinomas from the oral mucosa (Thomas et al, 1997), maximal $\beta 6$ expression in colon
20 cancer has been observed at the invading edges of tumour cell islands (Agrez et al, 1996).

Indeed, the $\beta 6$ subunit is widely observed in cancers of various origins (Breuss et al, 1995). For example, $\beta 6$ is detected in at least 50% of bowel cancer tumours. Others have reported its presence in oropharyngeal cancers where it is also present and strongly expressed in the invading margins of the cancer cell islands as is commonly found in
25 bowel cancer. In the oropharyngeal mucosa, no $\beta 6$ is observed in the normal lining cells of the mouth but in both primary and metastatic tumours from the oropharyngeal mucosa, strong $\beta 6$ expression is seen which does not correlate with degree of differentiation and in particular, is restricted to the basal layer of epithelial cells.

Expression of $\beta 6$ is also up-regulated in migrating keratinocytes at the wound edge
30 during experimental epidermal wound healing. $\alpha v \beta 6$ is not expressed in normal epithelium (Jones et al, 1997). However, following experimental wounding, αv appears to

switch its heterodimeric association from $\beta 5$ to $\beta 6$ subunit during re-epithelialisation. At day 3 after wounding, $\beta 6$ is absent but then appears around the perimeter of the basal cells of the migrating epidermis.

In human mucosal wounds, maximal expression of $\beta 6$ has been observed relatively late when epithelial sheets are fused and granulation tissue is present (Haapasalmi et al, 1996). Furthermore, those investigators observed maximal expression of tenascin with $\alpha v \beta 6$ expression. Interestingly, freshly isolated keratinocytes have not been found to express $\beta 6$ but begin to express this after subculturing. In contrast to persistent $\alpha v \beta 6$ expression observed in colon cancer cells, new expression of $\alpha v \beta 6$ in migrating keratinocytes is down-regulated to undetectable levels once re-epithelialisation is complete. However in normal unwounded skin, just as in other normal epithelia, $\alpha v \beta 6$ expression is absent indicating that this MAP kinase activation pathway is normally suppressed.

SUMMARY OF THE INVENTION

The present invention relates to the observation that the cellular activity of at least some types of cancer cells can be inhibited using an agent which binds to a binding domain of a MAP kinase for an integrin regardless of whether the integrin is expressed by the cells or not. It has also been found that some cancer cells may be substantially more susceptible to treatment with such an agent than corresponding normal cells. The reason for this is not known but this remarkable observation allows for essentially selective treatment of such cancers. In addition, the present invention in another form relates to the prophylaxis or treatment of blood cell cancers.

Accordingly, in one aspect of the present invention there is provided a method for prophylaxis or treatment of a cancer in a mammal, the method comprising treating the mammal with an effective amount of an agent that binds to a MAP kinase such that the capacity of the MAP kinase to bind with an integrin is inhibited, wherein the integrin is essentially not expressed by cancer cells of the cancer.

Typically, the agent will bind to a binding domain of the MAP kinase for the integrin.

Hence, in another aspect of the present invention there is provided a method for prophylaxis or treatment of a cancer in a mammal, the method comprising treating the mammal with an effective amount of an agent which binds to a binding domain of a MAP kinase for an integrin, wherein the integrin is essentially not expressed by cancer cells of
5 the cancer.

In a further aspect of the present invention there is provided a method for prophylaxis or treatment of a circulating blood cell cancer in a mammal, the method comprising treating the mammal with an effective amount of an agent which inhibits binding of the MAP kinase to the integrin.

10 In this instance, the agent may bind to either the MAP kinase or the integrin if the integrin is expressed by the cancer cells.

Typically, the agent will incorporate a carrier moiety for facilitating passage of the polypeptide across the cell membrane of the cancer cells. The carrier moiety will preferably be a signal peptide. The signal peptide will generally be a signal peptide for a
15 growth factor and most preferably, a signal peptide for Kaposi fibroblast growth factor (K-FGF). Usually, the signal peptide will comprise or consist of the amino acid sequence AAVALLPAVLLALLA, or a homologue, analogue, variant or derivative thereof.

Unexpectedly, it has been found that greater inhibition of cancer cell activity can be achieved with a signal peptide having or incorporating the amino acid sequence
20 AAVALLPAVLLALLA compared to other carrier moieties such as penetratin or for instance the $\beta 3$ signal peptide VTVLALGALAGVGVG. Preferably, the signal peptide will have the amino acid sequence AAVALLPAVLLALLAP. It is believed the greater degree of inhibition of cellular activity observed is due to the greater capacity of the signal peptide to translocate across the outer cellular membrane. This is surprising as it was
25 expected that the signal peptide for an integrin subunit chain would be more effective.

Accordingly, in another aspect of the present invention there is provided a method for prophylaxis or treatment of a cancer in a mammal, the method comprising administering to the mammal an effective amount of an agent incorporating a binding moiety which binds to a binding domain of a MAP kinase for an integrin and a signal
30 peptide having the amino acid sequence AAVALLPAVLLALLA for facilitating passage of

the binding moiety into cancer cells of the cancer, or a homologue, analogue, variant or derivative of the signal peptide.

In a further aspect of the present invention there is provided an agent for prophylaxis or treatment of a cancer in a mammal, the agent comprising a binding moiety which binds to a binding domain of a MAP kinase for an integrin and a signal peptide having the amino acid sequence AAVALLPAVLLALLA for facilitating passage of the binding moiety into cancer cells of the cancer, or a homologue, analogue, variant or derivative of the signal peptide.

Preferably, the agent utilised in a method of the invention will be a polypeptide. The polypeptide may for example be a fusion protein. Preferably, the polypeptide will comprise the binding domain of the integrin to which the MAP kinase binds. In the instance not all the amino acids in the binding domain are involved in the binding interaction with the MAP kinase, the polypeptide may comprise only those amino acids of the binding domain which are directly involved in the binding interaction.

Preferably, the polypeptide will comprise the amino acid sequence RSKAKWQTGTNPLYR, RARAKWDTANNPLYK, or RSRARYEMASNPLYR. Most preferably, the polypeptide will comprise an amino acid sequence selected from RSKAKNPLYR, RARAKNPLYK and RSRARNPLYR. Surprisingly, it has also been found that the amino acid sequence KEKLKNPLFK derived from the sequence KEKLKSQWNNDNPLFK of the cytoplasmic domain of the $\beta 2$ integrin subunit can bind to a MAP kinase.

Hence, in another aspect of the present invention there is provided a method for prophylaxis or treatment of a cancer in a mammal, the method comprising treating the mammal with an effective amount of an agent for inhibiting binding of the MAP kinase to a $\beta 2$ integrin subunit.

In yet another aspect of the present invention there is provided a method for modulating activity of cancer cells, the method comprising contacting the cells with an effective amount of a polypeptide having the sequence KEKLKNPLFK which binds to a binding domain of a MAP kinase for an integrin, or a homologue, analogue, variant or derivative thereof.

In still another aspect of the present invention there is provided an agent for modulating activity of cancer cells, the agent comprising a polypeptide having the sequence KEKLNPLFK, or a homologue, analogue, variant or derivative thereof.

There are also provided nucleic acid sequences encoding polypeptides,
5 homologues, analogues, and variants of the invention, recombinant vectors incorporating the nucleic acid sequences, and host cells transfected with such vectors. The vectors may for instance be cloning vectors, or expression vectors for expression of the polypeptides. In addition, the present invention relates to the use of the agents of the invention, including nucleic acids and vectors incorporating them, in the manufacture of
10 medicaments for the prophylaxis or treatment of cancer.

Typically, a polypeptide of the invention or administered to a mammal in accordance with the invention will have a length of about 150 amino acids or less, more preferably about 75 amino or 50 amino acids or less and most preferably, about 40 amino acids or less. When the polypeptide is a fusion protein or agent incorporating a carrier
15 moiety, the binding moiety that binds to the integrin will generally have a length of between about 5 to about 50 amino acids and more preferably, a length of between about 5 to about 35 amino acids.

Rather than administering a polypeptide to the mammal, a nucleic acid molecule encoding the polypeptide or a homologue or variant thereof may be administered for
20 expression of the polypeptide within the cancer cells. The nucleic acid sequence can be introduced into the cells in an appropriate expression vector for expression of the nucleic acid sequence extrachromosomally or more preferably, for integration of the nucleic acid sequence into genomic DNA by recombination events prior to expression of the polypeptide. Alternatively, the cells may be transfected with a nucleic acid molecule
25 incorporating nucleotide sequences flanking the sequence encoding the polypeptide which facilitate recombination with genomic DNA for expression of the polypeptide under the control of the transfected cell's own transcriptional regulatory sequences. Usually, the nucleic acid will be delivered intracellularly to the cancer cells by a viral vector. Adenovirus are preferred.

30 Preferably, the integrin will be a member of the αV integrin subfamily. More preferably, the integrin is or incorporates an integrin subunit selected from the group consisting of $\beta 2$, $\beta 3$, $\beta 5$, and $\beta 6$.

Preferably, the MAP kinase will be selected from the group consisting of an extracellular signal-regulated kinase (ERK), a JNK MAP kinase, and a p38 MAP kinase. Preferably, the MAP kinase is ERK1, ERK2 or JNK-1. Most preferably, the MAP kinase is ERK2.

5 By "cancer" is meant any type of unregulated cell proliferation. The cancer may for instance be selected from the group consisting of an epithelial cell cancer, prostate cancer, a lymphoma or blood cell cancers and other cancers, including leukemias such as myeloid leukemias, eosinophilic leukemias and granulocytic leukemias. Where the cancer is a blood cell cancer, the agent with which the mammal is treated may bind to an integrin
10 expressed by the blood cells of the cancer or to the MAP kinase such that binding of the MAP kinase with the integrin is thereby blocked or down-regulated. Typically, the agent will bind to the binding domain of the integrin for the MAP kinase or to the binding domain of the MAP kinase that binds to the integrin. In the instance the agent binds to the integrin, the agent will usually be a fragment of the MAP kinase comprising the
15 binding domain for the integrin, or a homologue, variant, analogue or derivative of such a fragment. Again, in the instance not all amino acids in the binding domain participate in the binding interaction, the agent may comprise only those amino acids of the binding domain which are directly involved in the binding interaction. The integrin may for example comprise $\beta 2$ the expression of which is restricted to white blood cells (Hines et al,
20 1992).

Suitable fragments of MAP kinases capable of binding with an integrin are described in International Patent Application No. WO 02/051993 and include the amino acid sequences HRDLKPSNLLLNTTCDLKICDFGLAR and PSNLLLNTTCDLKIC, and regions of such sequences.

25 The mammal may be any mammal treatable with a method of the invention. For instance, the mammal may be a member of the bovine, porcine, ovine or equine families, a laboratory test animal such as a mouse, rabbit, guinea pig, a cat or dog, or a primate or human being. Typically, the mammal will be a human being.

30 Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

In order that the nature of the present invention may be more clearly understood, preferred forms thereof will now be described with reference to the following non-limiting examples.

BRIEF DESCRIPTION OF THE FIGURES

5 **Figure 1** is a graph showing the binding of an ERK2 fragment to the cytoplasmic domain of $\beta 6$;

Figure 2 is a graph showing the binding of the MAP kinases ERK1 and ERK2 to a fragment comprising the binding domain of $\beta 6$;

10 **Figure 3** is a graph showing the binding of ERK2 to the peptide KEKLKNPLFK comprising the binding sequence of $\beta 2$;

Figure 4 is a graph showing the binding of the MAP kinases ERK2, p38 and JNK-1 to the cytoplasmic domain of $\beta 6$;

15 **Figure 5** is a graph showing the inhibition of the activity of HT29 colon cancer cells by a peptide comprising the binding domain of $\beta 6$ for ERK2 linked to the $\beta 3$ signal peptide or the modified K-FGF signal peptide AAVALLPAVLLALLA;

Figure 6 is a graph showing the inhibition of HT29 colon cancer cells using the peptide RSKAKNPLYR linked to penetratin or the modified K-FGF signal peptide AAVALLPAVLLALLA;

20 **Figure 7** is a graph showing a comparison of the inhibitory effect of the peptide RSKAKNPLYR linked to K-FGF signal peptide with and without a terminal proline residue on the proliferation of HT29 colon cancer cells;

Figure 8 shows FACScan analysis results for (A) normal human umbilical vein endothelial cells (HUVEC), (B) SH-SY5Y neuroblastoma cells, (C) HL60 leukemia cells, (D) DU 145 prostate cancer cells, and (E) HMEC-1 endothelial cells;

25 **Figure 9** shows FACScan analysis results for, (A) HT29 colon cancer cells, (B) H460 lung cancer cells and (C) melanoma cells;

Figures 10A to F are graphs showing the activity of the peptide RSKAKNPLYR linked to the modified K-FGF signal peptide AAVALLPAVLLALLA on different cell lines;

- 5 Figure 10G is a graph showing the activity of the AAVALLPAVLLALLARSARNPLYR peptide on HT29 colon cancer cells compared to the β 5 peptide RSRARYEMASNPLYR linked to the same signal peptide;

Figure 10H is a graph comparing the inhibitory effect of the peptide AAVALLPAVLLALLAPRSKAKNPLYR on the proliferation of HUVEC and HMEC-1 cells;

- 10 Figure 10I is a graph showing the inhibitory activity of the β 3 peptide RARAKNPLYK linked to AAVALLPAVLLALLA on the proliferation of HT29 colon cancer cells;

- 15 Figure 11 is a graph showing a comparison of the effect of the peptide AAVALLPAVLLALLAPRSKAKNPLYR compared to 5-fluorouracil on the proliferative response of HT29 colon cancer cells;

Figure 12 is a graph comparing the toxicity of the peptides AAVALLPAVLLALLARSKAKWQTGTNPLYR compared to 5-fluorouracil on HUVEC cells;

- 20 Figure 13 is a graph comparing the effect of the peptide AAVALLPAVLLALLAPRSKAKNPLYR and the chemotherapeutic agent CPT-11 on the proliferation of HT29 colon cancer cells;

Figures 14A and B are graphs showing the effect of the intra-tumoural injection of mice with the peptide AAVALLPAVLLALLARSKAKWQTGTNPLYR on the tumour growth of HT29 colon cancer cells and mouse weight, respectively; and

- 25 Figure 15 is a graph showing the effect of the intraperitoneal injection of mice with the peptide AAVALLPAVLLALLARSKAKWQTGTNPLYR on the tumour growth of HT29 colon cancer cells.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS OF THE PRESENT INVENTION

In the broadest sense, the term "integrin" unless otherwise specified, is to be taken to encompass an integrin family member or integrin subunit, or a homologue, derivative, variant or analogue of an integrin subunit, or an integrin family member incorporating at least one such homologue, derivative, variant or analogue of an integrin subunit.

By "binding domain" is meant the minimum length of contiguous amino acid sequence of the MAP kinase or integrin required for binding of the integrin or MAP kinase, respectively.

The term "homologue" is to be taken to mean a molecule that has amino acid sequence similarity. The homology between amino acid sequences is determined by comparing amino acids at each position in the sequences when optimally aligned for the purpose of comparison. The sequences are considered homologous at a position if the amino acids at that position are the same. Typically, a homologue will have an overall amino acid sequence homology of at least about 50% or 70% and most preferably, greater than about 80%, 90% or 98% sequence homology. Homology with a binding domain may be greater than the overall amino acid sequence homology of the homologue and will usually be greater than about 80% and preferably, greater than about 90%, 95% or 98%.

A homologue may be provided by, or the result of, the addition of one or more amino acids to an amino acid sequence, deletion of one or more amino acids from an amino acid sequence and/or the substitution of one or more amino acids with another amino acid or amino acids. Inversion of amino acids and other mutational changes that result in alteration of an amino acid sequence are also encompassed. A homologue may be prepared by introducing nucleotide changes in a nucleic acid sequence such that the desired amino acid changes are achieved upon expression of the mutagenised nucleic acid or for instance, by synthesising an amino acid sequence incorporating the desired amino acid changes.

The substitution of an amino acid may involve a conservative or non-conservative amino acid substitution. By conservative amino acid substitution is meant replacing an amino acid residue with another amino acid having similar stereochemical properties which does not substantially affect the conformation or the desired aspect or aspects of characteristic biological function. Preferred homologues include ones having amino acid

sequences in which one or more amino acids have been substituted with alanine or other neutrally charged amino acid residue(s), or to which one or more such amino acid residues have been added. A homologue may also incorporate an amino acid or amino acids not encoded by the genetic code.

- 5 By the term "variant" is meant an isoform of a polypeptide or fragment thereof, a naturally occurring mutant form of a polypeptide or fragment thereof, or a polypeptide or fragment thereof encoded by an allelic variant or partial nucleic acid sequence thereof.

- 10 The term "analogue" encompasses a molecule that differs from the original molecule but retains similarity in one or more features that provide the biological function characteristic of the original molecule. An analogue may have substantial overall structural similarity with the original molecule or only structural similarity with one or more regions of the original molecule responsible for the desired characteristic biological function. By "structural" similarity it is meant similarly in shape, conformation and/or other structural features responsible for the provision of the biological function or which otherwise have involvement in the provision of the biological function. Alternatively, it will be understood that with knowledge of the region(s) or domain(s) of a molecule having the desired characteristic biological function or knowledge of a binding domain to which the molecule binds, analogues may be provided that while differing in structure or chemical groups nevertheless possess such biological function. Indeed, an analogue may be a mimetic such as a peptido-mimetic. However, it is not necessary that an analogue have amino acid sequence homology, and an analogue may not be proteinaceous at all.

- 25 By the term "derivative" is meant a molecule that is derived or obtained from the original molecule and which retains one or more aspects or characteristic biological function of that molecule. A derivative may for instance be provided as a result of the cleavage of the original molecule, cyclisation and/or coupling with one or more additional moieties that improve solubility, lipophilic characteristics to enhance uptake by cells, stability or biological half-life, increased cellular toxicity, or for instance to act as a label for subsequent detection or the like. A derivative may also result from post-translational or post-synthesis modification such as the attachment of carbohydrate moieties, or chemical reaction(s) resulting in structural modification(s) such as the alkylation or acetylation of amino acid residues or other changes involving the formation of chemical bonds.

The term "polypeptide" is used interchangeably herein with "peptide" and encompasses amino acid sequences incorporating only a few amino acid residues or many amino acid residues coupled by peptide bonds. For instance, it will be understood that agents such as RSKAKNPLYR and KEKLNPLYR fall within the scope of the term.

5 The binding domain of an integrin to which a MAP kinase binds or the binding domain of the MAP kinase for the integrin may be identified and characterised using protocols and techniques described in International Patent Application No. WO 01/000677 and International Patent Application No. WO 02/051993, the disclosures of both of which are expressly incorporated herein by reference in their entirety.

10 More specifically, a binding domain may be localised by assessing the capacity of respective overlapping peptide fragments of the cytoplasmic domain of an integrin subunit or from a MAP kinase to bind with the MAP kinase or integrin, respectively. The specific amino acid sequence which constitutes the binding domain may then be determined utilising progressively smaller peptide fragments. In particular, test peptides
15 are readily synthesised to a desired length involving deletion of an amino acid or amino acids from one or both of the N-terminal and C-terminal ends of the larger amino acid sequence, and tested for their ability to bind with the MAP kinase or the integrin. This process is repeated until the minimum length peptide capable of binding with the MAP kinase or the integrin substantially without compromising the optimum observed level of
20 binding is identified.

 The identification of amino acids that play an active role in the MAP kinase integrin interaction may be achieved with the use of further synthesised test peptides in which one or more amino acids of the sequence are deleted or substituted with a different amino acid or amino acids to determine the effect on the ability of the peptide to bind with the
25 MAP kinase or the integrin. By deletion in this context is meant deletion of one or more of the amino acids between the N-terminal and C-terminal amino acid residues of the identified binding domain. Typically, substitution mutagenesis will involve substitution of selected ones of the amino acid sequence with alanine or other relatively neutrally charged amino acid.

30 Nucleotide and amino acid sequence data for the $\beta 6$ integrin subunit for instance is found in Sheppard et al, 1990. The nucleotide and amino acid sequence for ERK2 may be found in Boulton et al, 1991. Reference to such published data allows the ready design of

peptide fragments of an integrin subunit cytoplasmic domain for use in the identification and localisation of the binding domain of the integrin for the MAP kinase, and the identification of corresponding nucleic acid sequences encoding the peptide fragments.

Localisation and characterisation of a binding domain of an integrin for a MAP
5 kinase enables the design of agents which bind to the binding domain for modulation of cell activity. This will typically involve determining the physical properties of the binding domain such as size and charge distribution, and the tertiary structure of the binding domain. Specifically, at least the region of the integrin or MAP kinase containing the binding domain is modelled taking into account the stereochemistry and physical
10 properties of the binding domain such as size and charge distribution as well as its three dimensional structure as determined using x-ray crystallography, nuclear magnetic resonance and/or commercially available computer modelling software. Such modelling techniques are well known in the art. In a variation of this approach, the modelling will take into account the binding interaction of the binding domain with the MAP kinase or
15 the integrin such that any change in conformation arising from the interaction may be taken into account in the design of an analogue. Modelling flanking regions adjacent the binding domain also allows the design of agents for binding with such flanking regions but which are nevertheless capable of inhibiting the MAP kinase integrin interaction either by steric hindrance or by distorting the conformation of the binding domain of the
20 MAP kinase or integrin (eg. allosteric inhibitors).

The design of an analogue will usually involve selecting or deriving a template molecule onto which chemical groups are grafted to provide required physical and chemical characteristics or for further chemical reactions for achieving the required physical and chemical characteristics. The selection of template molecule and chemical
25 groups is based on ease of synthesis, likely pharmacological acceptability, risk of or potential for degradation *in vivo*, stability and maintenance of biological activity upon administration.

In order to constrain a polypeptide or other agent in a three dimensional conformation required for binding, it may be synthesised with side chain structures or be
30 synthesised incorporating cysteine residues which form a disulfide bridge, or otherwise be incorporated into a molecule with a known stable structure *in vivo*. For example, a

polypeptide or the like may be incorporated into an amino acid sequence at least part of which folds into a β -pleated sheet or helical structure such as an α -helix.

5 A polypeptide or other agent may also be cyclised to provide enhanced rigidity and thereby stability *in vivo*. Various methods for cyclising peptides, fusion proteins or the like are known (eg. Schiller et al., 1985). For example, a synthetic peptide incorporating two cysteine residues distanced from each other along the peptide may be cyclised by the oxidation of the thiol groups of the residues to form a disulfide bridge between them. Cyclisation may also be achieved by the formation of a peptide bond between the N-terminal and C-terminal amino acids of a synthetic peptide or for instance through the
10 formation of a bond between the positively charged amino group on the side chain of a lysine residue and the negatively charged carboxyl group on the side chain of a glutamine acid residue. As will be understood, the position of the various amino acid residues between which such bonds are formed will determine the size of the cycle. Variation of cycle size for optimisation of binding affinity may be achieved by synthesising peptides in
15 which the position of amino acids for achieving cyclisation has been altered. The formation of direct chemical bonds between amino acids or the use of any suitable linker to achieve a desired three-dimensional conformation is also well within the scope of the skilled addressee.

Strategies for identifying agents suitable for use in methods of the present invention
20 include large scale screening techniques. For example, peptide library protocols provide an efficient way of testing a vast number of potential agents. Such libraries and their use are well known. Prospective agents identified may be then further evaluated in suitable activity, competitive and other assays. A method of screening for an agent or evaluating whether an agent is capable of binding to the binding domain of an integrin for a MAP
25 kinase and thereby inhibiting the MAP kinase integrin interaction, may for instance involve utilising the agent in an assay whereby the agent has the opportunity of binding to the MAP kinase in the presence of the integrin prior to the addition of the integrin, and determining whether inhibition of binding of the MAP kinase to the integrin results. An alternate screening method may for instance involve selecting a test agent and measuring
30 cellular activity of target cells in the presence of the test agent, and comparing that activity with cellular activity in the absence of the test agent. Cellular activity may be assessed by cell growth as indicated by [^3H]-thymidine uptake or other measurement of

cellular activity. As will be understood, a difference in observed functional activity in the presence of the test agent is indicative of the modulating effect provided by the test agent.

5 It will be understood that the integrin in the context of such assays may be an integrin subunit or polypeptide or fragment incorporating the binding domain of the integrin to which the MAP kinase binds, or a homolog, analog, variant or derivative of such a molecule to which the MAP kinase is capable of binding. Similarly, a MAP kinase in this context may be an intact MAP kinase or a fragment thereof incorporating a binding domain for the integrin, or a homolog, analog, variant or derivative thereof that is capable of binding with the integrin.

10 Determination of whether an agent is capable of binding to the binding domain of the MAP kinase or integrin may be achieved using a polypeptide or fragment as described herein consisting of the binding domain of the integrin or MAP kinase or the binding sequence or fragment thereof of the binding domain that directly participates in the binding interaction, or analogs or the like of such molecules.

15 In this regard, the regions or amino acids in the binding domain that participate in the binding interaction may be separated by amino acids that are not involved in the binding interaction. The binding sequence of a binding domain is the sequence of amino acids which participate in the binding interaction excluding those amino acids in the binding domain that do not participate in the binding interaction. As an example, the
20 binding domain of $\beta 6$ comprises the amino acid sequence RSKAKWQTGTNPLYR. However, the intervening amino acid sequence WQTGT does not directly participate in binding with the MAP kinase ERK2. That is, even if the sequence WQTGT is deleted, a peptide with the amino acid sequence RSKAKNPLYR is still bound by ERK2. Similarly, the binding domains of $\beta 3$ and $\beta 5$ for ERK2 comprise RARAKWDTANNPLYK and
25 RSRARYEMASNPLYR, respectively. Deletion of the intervening sequences WDTAN and YEMAS from the sequences yields the peptides RARAKNPLYK and RSRARNPLYR both of which are still bound by ERK2.

Alignment of binding domains of $\beta 2$, $\beta 3$ and $\beta 5$ and $\beta 6$ results in the consensus
30 scheme R/K x R/K * R/K - xx*x* NPL Y/F R/K wherein R/K is either arginine or lysine, Y/F is either tyrosine or phenylalanine, x may be any amino acid, * is a hydrophobic amino acid residue, and - is an amino acid (serine) present in the binding

domain of $\beta 2$ but not in the others and which may be replaced with another amino acid such as threonine, tyrosine, asparagine or glutamine, or be deleted. Accordingly, peptides are provided that comprise or consist of this sequence that may find application in methods of the invention. As indicated above, the intervening amino acid sequence indicated by $-xx*x*$ may also be deleted such that an agent of the invention comprising or consisting of the sequence R/KxR/K*R/KNPLY/FR/K is provided.

A particularly preferred way of achieving intracellular delivery of polypeptides, nucleic acids and other agents is to use "carrier peptides" which have the ability to deliver macro-molecules across cell membranes in an energy-independent manner. Indeed, such peptides provide the possibility of both testing potential agents in cell culture without drastically altering cell membrane integrity and of delivering agents *in vivo*. Carrier peptides that are known in the art include penetratins and variants thereof (Derossi et al, 1994, 1996), human immunodeficiency virus Tat derived peptide (Prociantz, 1996), transportan derived peptide (Pooga et al, 1998) and signal peptides.

Particularly preferred signal peptides are described in United States Patent No. 5807,746 the contents of which are incorporated herein in its entirety. In particular, a signal peptide for Kaposi fibroblast growth factor (K-FGF) having or incorporating the amino acid sequence AAVALLPAVLLALLA or AAVALLPAVLLALLAP is preferred.

It is not necessary that the signal peptide used in a method of the invention be a complete signal peptide and fragments or homologues and the like which retain the ability to penetrate the outer cellular membrane to effect delivery of the attached agent into the cytosol of the cell may be utilised. In the instance the agent is a nucleic acid, the signal peptide will typically also be capable of penetrating through the nuclear membrane of eukaryotic cells and thereby effecting translocation of the attached nucleic acid into the nucleus of the cell.

The agent may be linked to the carrier peptide in any conventionally known manner. For instance, a polypeptide may be linked directly to a carrier peptide through an amino acid linker sequence by a peptide bond or non-peptide covalent bond using a crosslinking reagent. For agents that have a negative charge such as nucleic acids, the agent may be linked to the carrier peptide by charge-association between the negatively charged agent and the positively charged amino acids in the carrier peptide or linker

sequence. Chemical ligation methods may also be used to create a covalent bond between the carboxy terminal amino acid of the carrier peptide or linker sequence and the agent.

Specific targeting of agents to abnormal cells can be realised by coupling a ligand or antibody or binding fragment thereof (such as *Fab* and *F(ab)₂* fragments) to carrier molecules such as penetratin which are in turn coupled to the agent. Another approach may employ coupling the carrier-agent complex to integrin receptor-targeted peptides which target an extracellular integrin domain. For example, peptides with the sequence DLXXL can be used to target the extracellular domain of $\beta 6$. Given that $\beta 6$ expression enhances effective proteolysis at the cell surface by matrix metalloproteinase-9 (MMP-9) (Agrez et al, 1999), such targeting approaches include engineering an MMP-9 cleavage site between the targeting moiety and the carrier to facilitate internalisation of the carrier-agent complex. As another example, the ligand recognition motif for $\alpha V\beta 6$ integrin, RTDLSLRITYTL may be used in conjunction with or without an engineered MMP-9 cleavage site to deliver the carrier-agent complex to the surface of the target cell. Further protocols for delivering nucleic acids to cells by targeting integrins is described in Bachmann et al, 1998.

The provision of fusion proteins and use of fusion proteins incorporating a polypeptide which binds to the binding domain of a MAP kinase for an integrin is expressly provided for by the invention. Polypeptides and fusion proteins or the like may be synthesised or produced using conventional recombinant techniques. Nucleic acid encoding a fusion protein may for instance be provided by joining separate DNA fragments encoding peptides or polypeptides having desired three dimensional conformations and/or amino acid sequences by employing blunt-ended termini and oligonucleotide linkers, digestion to provide staggered termini as appropriate, and ligation of cohesive ends. Alternatively, PCR amplification of DNA fragments can be utilised employing primers which give rise to amplicons with complementary termini which can be subsequently ligated together (eg. see Current Protocols in Molecular Biology. John Wiley & Sons, 1992).

Polypeptides and fusion proteins may be expressed *in vitro* and purified from cell culture for administration to a subject, or cells may be transfected with nucleic acid encoding a polypeptide or fusion protein for *in vitro* or *in vivo* expression thereof. The nucleic acid will typically first be introduced into a cloning vector and amplified in host

cells, prior to the nucleic acid being excised and incorporated into a suitable expression vector for transfection of cells.

Typical cloning vectors incorporate an origin of replication (*ori*) for permitting efficient replication of the vector, a reporter or marker gene for enabling selection of host cells transformed with the vector, and restriction enzyme cleavage sites for facilitating the insertion and subsequent excision of the nucleic acid sequence of interest. Preferably, the cloning vector has a polylinker sequence incorporating an array of restriction sites. The marker gene may be drug-resistance gene (eg. Amp^r for ampicillin resistance), a gene encoding an enzyme such as chloramphenicol acetyltransferase (CAT), β -lactamase, adenosine deaminase (ADA), aminoglycoside phosphotransferase (APH), dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), or for instance β -galactosidase encoded by the *E. coli lacZ* gene (*LacZ*). Yeast reporter genes include imidazole glycerolphosphate dehydratase (HIS3), *N*-(5'-phosphoribosyl)-anthranilate isomerase (TRP1) and β -isopropylmalate dehydrogenase (LEU2). As will be appreciated, expression vectors of the invention may also incorporate such marker genes.

Cloning vectors include cloning vectors for mammalian, yeast and insect cells. Particular vectors that may find application include pBR322 based vectors and pUC vectors such as pUC118 and pUC119. Suitable expression and cloning vectors are for instance described in Molecular Cloning. A Laboratory Manual., Sambrook et al., 2nd Ed. Cold Spring Harbour Laboratory., 1989.

Suitable expression vectors include plasmids and cosmids capable of expression of a DNA (eg. genomic DNA or cDNA) insert. An expression vector will typically include transcriptional regulatory control sequences to which the inserted nucleic acid sequence is operably linked. By "operably linked" is meant the nucleic acid insert is linked to the transcriptional regulatory control sequences for permitting transcription of the inserted sequence without a shift in the reading frame of the insert. Such transcriptional regulatory control sequences include promoters for facilitating binding of RNA polymerase to initiate transcription, expression control elements for enabling binding of ribosomes to transcribed mRNA, and enhancers for modulating promoter activity. A promoter may be a tissue specific promoter which facilitates transcription of the nucleic acid insert only in specific cell lineages and not in other cell types or only to a relatively low level in such other cell types. The design of an expression vector will depend on the

host cell to be transfected, the mode of transfection and the desired level of transcription of the nucleic acid insert.

Numerous expression vectors suitable for transfection of prokaryotic (eg. bacterial) or eukaryotic (eg. yeast, insect or mammalian cells) are known in the art. Expression vectors suitable for transfection of eukaryotic cells include pSV2neo, pEF.PGK.puro, pTk2, pRc/CNV, pcDNA1/neo, non-replicating adenoviral shuttle vectors incorporating the polyadenylation site and elongation factor 1- α promotor and pAdEasy based expression vectors most preferably incorporating a cytomegalovirus (CMV) promotor (eg. see He et al, 1998). For expression in insect cells, baculovirus expression vectors may be utilised examples of which include pVL based vectors such as pVL1392, and pVL941, and pAcUW based vectors such as pAcUW1. Viral expression vectors are particularly preferred.

Any means for achieving the introduction of the nucleic acid into a target cell may be used. Transfer methods known in the art include viral and non-viral transfer methods. Suitable virus into which appropriate viral expression vectors may be packaged for delivery to target cells include adenovirus, vaccinia virus, retroviruses of avian, murine and human origin, herpes viruses including Herpes Simplex Virus (HSV) and EBV, papovaviruses such as SV40, and adeno-associated virus. Particularly preferred virus are replication deficient recombinant adenovirus (eg. He et al, 1998). Engineered virus may be administered locally or systemically to achieve delivery of nucleic acid sequence into a target cell.

Rather than utilising viral mediated transfection of cells, nucleic acid sequences and other agents may be introduced into a cell *in vitro* or *in vivo* by liposome mediated transfection. The liposomes may carry targeting molecules for maximising delivery of the agent or agents contained therein to specific cell types of interest. Such targeting molecules include antibodies or binding fragments thereof as described above, ligands or cell surface receptors for facilitating fusion of liposomes to the specific cells of interest. Agents may also be intracellularly delivered *in vitro* using conventional cold or heat shock techniques or for instance, calcium phosphate coprecipitation or electroporation protocols as are known in the art. Yet another strategy is to design the agent to have the inherent ability to pass across the lipid bilayer of a cell.

Host cells that may be used for expression of polypeptides or fusion proteins include bacteria such as *E. coli*, Bacillus such as *B. subtilis*, *Streptomyces* and *Pseudomonas* bacterial strains, yeast such as *Sacchromyces* and *Pichia*, insect cells, avian cells and mammalian cells such as Chinese Hamster Ovary cells (CHO), COS, HeLa, HaRas, WI38, SW480, and NIH3T3 cells. The host cells are cultured in a suitable culture medium under conditions for facilitating expression of the introduced nucleic acid prior to purification of the expressed product from the host cells, and/or supernatants as the case may be using standard purification techniques.

The toxicity profile of an agent may be tested on normal and abnormal cells such as cancer cells by evaluation of cell morphology, trypan-blue exclusion, assessment of apoptosis and cell proliferation studies (eg. cell counts, ³H-thymidine uptake and MTT assay).

Agents of the invention may be co-administered with one or more other compounds or drugs. For example, an agent or agents may be co-administered in combination or in conjunction with antisense therapy or chemotherapeutic drugs. By "co-administered" is meant simultaneous administration in the same formulation or in two different formulations by the same or different routes, or sequential administration by the same or different routes. By "sequential" administration is meant one is administered after the other, typically with a time delay of from very short times up to hours or for instance days.

The agent or agents will typically be formulated into a pharmaceutical composition incorporating a pharmaceutically acceptable carriers and/or excipient for administration to the intended subject. Pharmaceutical compositions include sterile aqueous solutions suitable for injection, (where the agent or agents is water soluble) and sterile powders for the extemporaneous preparation of sterile injectable solutions. Such injectable compositions will be fluid to the extent that the syringability exists and typically, will be stable to allow for storage after manufacture. The carrier may be a solvent or dispersion medium containing one or more of ethanol, polyol (eg. glycerol, propylene glycol, liquid polyethylene glycol and the like), vegetable oils and mixtures thereof. Fluidity may be maintained by the use of a coating such as lecithin and by the use of surfactants.

Sterile injectable solutions will typically be prepared by incorporating the active agents in the desired amount in the selected solvent with various other components

enumerated above, prior to sterilising the solution by filtration. Generally, dispersions will be prepared by incorporating the sterile active agents into a sterile vehicle which contains the dispersion medium and other components. In the case of sterile powders, preferred methods of preparation are vacuum drying and freeze-drying techniques which yield a powder of the active agent plus any additional desired ingredient from previously sterile filtered solutions thereof.

For oral administration, the active agents may be formulated into any orally acceptable carrier deemed suitable. In particular, the active ingredient may be formulated with an inert diluent, an assimilable edible carrier or it may be enclosed in a hard or soft shell gelatin capsule. Moreover, an active agent may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions or syrups.

Active agents may also be formulated into topically acceptable carriers conventionally used for forming creams, lotions or ointments for internal or external application. Topical formulations may be applied to a site to be treated by dressings and the like impregnated with the formulation.

Typically, a composition of the invention will incorporate one or more preservatives such as parabens, chlorobutanol, phenol, sorbic acid, and thimersal. In many cases, a composition may furthermore include isotonic agents such as sugars or sodium chloride. In addition, prolonged absorption of the composition may be brought about by the use in the compositions of agents for delaying absorption such as aluminium monostearate and gelatin.

Tablets, troches, pills, capsules and the like may also contain one or more of the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; a disintegrating agent such as corn starch, potato starch or alginic acid; a lubricant such as magnesium stearate; a sweetening agent such as sucrose, lactose or saccharin; and a flavouring agent.

Pharmaceutically acceptable carriers include any suitable conventionally known solvents, dispersion media and isotonic preparations or solutions. Use of such ingredients and media for pharmaceutically active substances is well known. Except insofar as any conventional media or agent is incompatible with the active agent, use

thereof in therapeutic and prophylactic compositions is included. Supplementary active ingredients can also be incorporated into the compositions if desired.

It is particularly preferred to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein is to be taken to mean physically discrete units suited as unitary dosages for the subject to be treated, each unit containing a predetermined quantity of active agent calculated to produce the desired therapeutic or prophylactic effect in association with the relevant carrier used.

When the dosage unit form is a capsule, it may contain in addition to one or more of the above ingredients a liquid carrier. Various other ingredients may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills or capsules may be coated with shellac, sugars or both. In addition, an active agent may be incorporated into any suitable sustained-release preparation or formulation.

Pharmaceutical compositions will generally contain at least about 1% by weight of the active agent or agents. The percentage may of course be varied and may conveniently be between about 5 to about 80% w/w of the composition or preparation. As will be appreciated, the amount of active agent or agents in such compositions will be such that a suitable effective dosage will be delivered to the subject taking into account the proposed mode of administration. Preferred oral compositions according to the invention will contain between about 0.1 μ g and 4000mg of the active agent.

In addition, a pharmaceutical composition may contain a vector of the invention capable of transfecting target cells. The vector may for instance, be packaged into a suitable virus for delivery of the vector into target cells as described above.

The dosage of an active agent will depend on a number of factors including whether the agent is to be administered for prophylactic or therapeutic use, the condition for which the agent is intended to be administered, the severity of the condition, the age of the subject, and related factors including weight and general health of the subject as may be determined by the physician or attendant in accordance with accepted principles. For instance, a low dosage may initially be given which is subsequently increased at each administration following evaluation of the subjects response. Similarly, frequency of administration may be determined in the same way that is, by continuously monitoring

the subject's response between each dosage and if necessary, increasing the frequency of administration or alternatively, reducing the frequency of administration. Typically, an agent will be administered in accordance with a method of the invention at a dosage up to about 50mg/kg body weight of the mammal and preferably in a range of from about
5 20mg/kg to 40mg/kg body weight.

The route of administration of a pharmaceutical composition will again depend on the nature of the condition for which the composition is to be administered. Suitable routes of administration include but are not limited to respiratorily, intratracheally, nasopharyngeally, intravenously, intraperitoneally, subcutaneously, intracranially,
10 intradermally, intramuscularly, intraocularly, intrathecally, intranasally, by infusion, orally, rectally, via IV group patch, topically and by implant. With respect to intravenous routes, particularly suitable routes are via injection into blood vessels which supply a tumour or particular organs to be treated. Agents may also be delivered into cavities such for example the pleural or peritoneal cavity, or be injected directly into tumour tissue.

15 The present invention will be described herein after with reference to a number of examples.

Example 1: Binding of MAP kinases to the cytoplasmic domain of integrin subunit chains.

1.1: ELISA assay for detection of MAP kinase binding

20 An assay was developed for detecting the ability of MAP kinases to bind with a peptide agent in solution.

Briefly, the MAP kinase is added to an eppendorf tube containing a biotinylated test peptide in phosphate buffered saline (PBS) at pH 7.4. The mixture is allowed to stand at room temperature for 10 minutes with occasional mixing by gentle inversion of the tube.
25 The MAP kinase/test peptide mixture is then transferred to a pre-washed (PBS x 3 x 100 µl) 96-well streptavidin coated ELISA microtitre plate and allowed to stand for 60 minutes at room temperature. At the end of the 60 minute period, the MAP kinase/test peptide mixture is tipped from the ELISA plate and the wells washed 3 times with wash solution (PBS + 0.05% Tween - 20).

Primary antibody detection for the MAP kinase is then added to the test wells and the ELISA plate is allowed to stand for a further 30 minutes at room temperature before the antibody solution is decanted from the plate and the wells again washed with the wash solution. Optimal dilutions of primary detection antibody are pre-determined for each target MAP kinase. Alkaline-phosphatase or other enzyme conjugated anti-primary species antibody at a dilution of 1:2000 v/v is subsequently added to the wells and the microtitre plate is allowed to stand for another 30 minutes at room temperature, before the washing step is repeated and phosphatase detection reagent is added to the wells and the colour reaction allowed to develop. The wells are read at 405 nm using a microplate reader (Bio-Rad). Controls comprise non-biotinylated peptide and biotinylated peptide alone

1.2: ERK, JNK and p38 MAP kinases bind to integrins

The binding of various MAP kinases to the cytoplasmic domain of $\beta 6$ was tested using the assay described in Example 1.1. Specifically, the biotinylated ERK2 amino acid sequence biotin-HRDLKPSNLLLNTTCDLKICDFGLAR was incubated with the cytoplasmic domain of $\beta 6$ having the amino acid sequence HDRKEVAKFEAERSKAKWQTGTNPLYRGSTSTFKNVTYKHREKQKVDLSTDC, and the binding interaction detected using the anti- $\beta 6$ antibody R6G9. The ERK fragment bound to the $\beta 6$ domain is shown in Fig. 1.

Similarly, the binding of ERK1 and ERK2 to the fragment RSKAKWQTGTNPLYR comprising the binding domain of the $\beta 6$ integrin subunit was tested using the assay of Example 1.1 and the results are shown in Fig. 2. As can be seen, both ERK1 and ERK2 bind to the $\beta 6$ fragment. ERK1 and ERK2 used for all studies was obtained by purification of the protein following cleavage of GST-ERK1/2 with thrombin. GST-ERK1/2 are fusion proteins consisting of ERK coupled to glutathione-S-transferase and purified from host cells transfected with pGEX-4T vector.

The fragment of $\beta 2$ corresponding to the amino acid sequence RSKAKWQTGTNPLYR of $\beta 6$ is KEKLKSQWNNDNPLFK. The peptide RSKAKNPLYR (arbitrarily designated peptide 10(4)) obtained by the deletion of the intervening sequence WQTGT from the $\beta 6$ fragment also binds to ERK2 (see Example)). To test whether deletion of SQWNND from the $\beta 2$ fragment had any effect, the peptide KEKLKNPLFK

(designated $\beta 2$ (10(4))) was synthesised and incubated with ERK2. As shown in Fig. 3, the peptide was bound by ERK2.

Next, fragments of ERK2, p38 MAP kinase and JNK-1 MAP kinases were tested for ability to bind with the cytoplasmic domain fragment of $\beta 6$

- 5 HDRKEVAKFEAERSKAKWQTGTNPLYRGSTSTFKNVITYKHREKQKVDLSTDC. The amino acid sequence for the ERK2 fragment used is HRDLKPSNLLLNTTCDLKICDFGLAR while that for the p38 MAP kinase and JNK-1 MAP kinase fragments are HRDLKPSNLAVNEDCELKILDFGLAR and HRDLKPSNIVVKSDCTLKILDFGLAR, respectively . The results are shown in Fig. 4. All
10 of the MAP kinase fragments bound to the $\beta 6$ cytoplasmic domain.

Example 2: Effect of the carrier moiety on inhibition of HT29 cancer cell activity

2.1: Cell proliferation assay using MTT

- An assay using the substrate methylthiazolotetrazolium (MTT) was utilised to determine the inhibitory affect of peptide agents on cancer cell proliferation. MTT
15 substrate (CAT No. M-2128, Sigma) is cleaved in growing cells to yield a water insoluble salt. After solubilisation of the salt, a coloured product is produced that allows quantitation of the proliferative activity of the cultured cells. To prepare the MTT solution for use in the assay, 100 mg of MTT is mixed with 20 ml of PBS at pH 7.4. The resulting solution is filter sterilised (0.2 μ M syringe filter) and stored at 4°C protected
20 from light until use.

- Cells from the colon cancer cell line HT29 were seeded into wells of a 96-well microtitre plate in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% foetal bovine serum, glutamine, Hepes, and antibiotics at a concentration of 2,000 cells/100 μ l and cultured for 24 hours in 5% CO₂ in air at 37°C. After the 24 hour
25 incubation period the test agent (solubilised in complete DMEM) is added to test wells to a final media volume of 200 μ l and the microtitre plate is incubated for a further 48 to 72 hour period.

- A 20 μ l volume of the MTT solution prepared as above is then added to the test wells and the microtitre plate incubated for further 3 hours in 5% CO₂ in air at 37°C. The
30 plate is subsequently centrifuged for 5 minutes at 2,000 rpm in a microplate carrier. The

supernatant is then removed from the wells using a multichannel pipette leaving MTT crystals in the wells.

To the wells is then added 150 μ l of 6:1 v/v DMSO/glycine solution (0.1 M glycine and 0.1 M NaCl adjusted to pH 10.5 using NaOH) and the MTT crystals are dissolved by gently vortexing the microtitre plate. Absorbance is read at 550 nm using a microtitre plate reader. The percentage inhibition of proliferation of the test cells is calculated relative to untreated control cells.

2.2: Synthesis and testing of peptide agents

Cells from the colon cancer cell line HT29 express β 6. To assess the effect of carrier moieties on inhibition of cellular activity of HT29 cells by peptides which bind to ERK2, peptides comprising the β 6 fragment RSKAKWQTGTNPLYR linked directly to the β 3 signal peptide VTVLALGALAGVGVG (Liu et al, PNAS (1996) 93: 11819-11824) or AAVALLPAVLLALLA (designated nfkb) in which the terminal proline of the K-FGF signal peptide AAVALLPAVLLALLAP (designated nfkbp) (US Patent No. 6,248,558) has been deleted were synthesised as follows:

VTVLALGALAGVGGRSKAKWQTGTNPLYR (Frag 5B3)

AAVALLPAVLLALLARSKAKWQTGTNPLYR (Frag 5 nfkb)

The inhibition of the activity of the HT29 cells by the synthetic peptides was assessed using the cell proliferation assay described in Example 2.1 and the results are shown in Fig. 5. Use of the modified K-FGF signal peptide resulted in total inhibition of the proliferation of the HT29 colon cancer cells at the higher concentration of the peptides used. In contrast, use of the β 3 signal peptide resulted in a decrease in proliferation of the cells to only about 37% of that of the control cells.

Synthetic peptides comprising the 10(4) peptide RSKAKNPLYR linked directly to penetratin (RQIKWIFQNRRMKWKKC) or the modified K-FGF signal peptide AAVALLPAVLLALLA were also prepared and their inhibitory effect of the proliferation of the HT29 colon cancer cells assessed as described above. The complete sequences for these peptides are as below and the results are shown in Fig. 6.

AAVALLPAVLLALLARSKAKNPLYR (Ik-2 (sig))

RQIKIWFQNRRMKWKKRSKAKNPLYR (Ik-2 (pen))

A greater degree of inhibition was observed using the modified K-FGF signal peptide at all concentrations of the synthetic peptides tested.

5 A comparison of the inhibitory effect of the 10(4) peptide RSKAKNPLYR linked to the K-FGF signal peptide with and without the terminal proline residue on the proliferation of the HT29 cells is shown in the Fig. 7. The complete sequences for the peptides are as follows:

AAVALLPAVLLALLARSKAKNPLYR (nfkb10(4)acetate)

AAVALLPAVLLALLAPRSKAKNPLYR (nfkb10(4)p)

10 As indicated in Fig.7, the presence of the terminal proline of the K-FGF signal peptide resulted in a greater degree of inhibition of proliferation of the HT29 cells over the concentrations of the peptides used with total inhibition being achieved by both synthetic peptides at the highest concentration employed.

Example 3: Inhibition of $\beta 6$ expressing and non-expressing cells.

15 3.1 FACScan Analysis

Different cell lines were assessed for $\beta 3$, $\beta 5$ and $\beta 6$ expression by FACScan analysis. Briefly, monolayer cultures of the cells were harvested with trypsin/EDTA. Cells were washed once with PBS, incubated with primary antibody against integrin subunits for 20 minutes at 4°C and then washed twice with PBS. Cells were then
20 stained with secondary antibody conjugated with phycoerythrin for 20 minutes at 4°C, washed twice with PBS and resuspended in 0.5 ml PBS prior to FACScan analysis (Becton Dickenson, Rutherford, New Jersey, USA).

FACScan results for normal human umbilical vein endothelial cells (HUVEC), the neuroblastoma cell line SH-SY5Y, the leukemia cell line HL 60, the prostate cancer cell
25 line was DU 145 and the HMEC-1 cell line are shown in Fig. 8A to E, respectively. HMEC-1 cells are malignant HUVEC cells which have been transformed with the virus SV40. As indicated in the graphs shown, all of these cell lines are essentially non- $\beta 6$ expressing cells but do express $\beta 3$ and $\beta 5$ with the exception of the SH-SY5Y cell line which is also essentially non- $\beta 5$ expressing. In contrast, the colon cancer cell line HT29,

the lung cancer cell line H460, and the melanoma cell line WM 115 are all $\beta 6$ expressing cells as shown in Fig. 9A to C, respectively. The breast cancer cell line MCF-7 is also $\beta 6$ expressing (data not shown).

3.2 Inhibition of cell proliferation

5 The ability of the 10(4) peptide RSKAKNPLYR linked to the signal peptide AAVALLPAVLLALLA to inhibit the proliferation of the above cell lines was assessed using the cell proliferation assay described in Example 2.1.

10 As shown in Fig. 10A, administration of the peptide agent to HUVEC cells has essentially no impact on the proliferation of cells. In contrast, administration of the peptide agent to the leukaemia cell line HL60, the prostate cancer cell line DU145 and the HMEC-1 cell line resulted in increasing levels of inhibition of cellular activity as concentrations of the peptide increased with total inhibition of proliferation being observed at the highest concentration of the peptide as indicated in Fig. 10B to Fig. 10D. The same result was found for the lung cancer cell line H460 and the melanoma cell line 15 WM115 using the peptide agent as shown in Fig. 10E and Fig. 10F, respectively. The inhibition of proliferation of HT29 colon cancer cells when treated with the $\beta 5$ integrin-derived RSRARYEMASNPLYR peptide or RSRARNPLYR linked with the modified signal peptide AAVALLPAVLLALLA (designated nf-b5-frag 5 and nf-b5-10-4, respectively) is shown in Fig. 10G. While both peptides inhibited proliferation of the 20 cells, a greater degree of inhibition was observed for the peptide agent incorporating RSRARNPLYR.

25 A comparison of the inhibitory effect of the peptide AAVALLPAVLLALLAPRSKAKNPLYR (nfkb10(4)p) on the proliferation of HUVEC and HMEC-1 cells shown in Fig. 10H. As can be seen, increasing inhibition of proliferation of the HMEC-1 cell line was observed with increasing concentration of the peptide while inhibition of proliferation of the HUVEC cells was relatively minimal.

The finding that the malignant HMEC-1 cells but not normal HUVEC cells are inhibited by the peptide agent shows selective activity of the peptide agent on the cancer cells.

Inhibition of proliferation of HT29 cells using a peptide having the amino acid sequence RARAKWDTANNPLYK comprising the binding domain of $\beta 3$ (designated nfkb- $\beta 3$) for ERK2 is shown in Fig. 10I. The signal peptides AAVALLPAVLLALLA and AAVALLPAVLLALLAP alone have essentially no effect on the proliferation of any of the cell lines tested (data not shown).

Example 4: Comparison with conventional anti-cancer drugs

4.1: Effect of $\beta 6$ cytoplasmic derived peptide on HT29 colon cancer cell proliferation compared to 5-fluorouracil (5-FU)

HT29 colon cancer cells were treated with either the anti-cancer drug 5-FU or the 10(4) peptide RSKAKNPLYR linked to the K-FGF signal peptide AAVALLPAVLLALLAP. As can be seen in Fig. 11, the peptide (designated nfkb10(4)p) was more effective at inhibiting the proliferative response of the HT29 colon cancer cells compared to 5-FU.

The effect of the AAVALLPAVLLALLARSKAKWQTGTNPLYR peptide agent on normal HUVEC cells compared to 5-FU is shown in Fig. 12. The peptide agent is designated IK-1 in the figure. The graph shows that the peptide has essentially no impact on the cells in contrast to the toxicity of the chemotherapeutic agent 5-FU. Fig. 13 further shows that the 10(4) peptide RSKAKNPLYR linked directly to the K-FGF signal peptide AAVALLPAVLLALLAP (designated nfkb10(4)p in the figure) is more effective in inhibiting proliferation of the colon cancer cell line HT29 compared to the chemotherapeutic agent CPT-11 at all concentrations used.

Example 5: Effect of peptide agents on *in vivo* tumour growth in Balb/c Nu/Nu nude mice

The ability of the peptide agent AAVALLPAVLLALLARSKAKWQTGTNPLYR (designated IK-1) to inhibit tumour growth in immune-deficient mice was assessed.

Balb/c female athymic mice (8 weeks of age obtained from the Animal Resource Centre, Perth, Western Australia) were maintained under pathogen-free conditions and fed standard mouse chow and water *ad lib*. The mice were divided into groups of ten each. Mice in different groups were marked by ear notching. Briefly, the mice were anaesthetised using isoflurothane by inhalation and their ears marked using a surgical

punch. After a recovery period of 48 hours, the mice in each group were injected with 1×10^6 HT29 colon cancer cells in standard DMEM culture medium subcutaneously dorsally behind the right shoulder using a 21-gauge needle. Animal weights and tumour sizes (breadth and length measured with callipers) were recorded weekly. Mice were sacrificed by CO₂ asphyxiation and tumour mass excised for measurement.

Mice were injected twice weekly for four weeks commencing 10 days after tumour cell innoculation with DMEM, signal peptide (AAVALLPAVLLALLA) (designated "carrier") or the IK1 AAVALLPAVLLALLARSKAKWQTGTNPLYR peptide agent, and tumour growth was compared to an untreated control group.

Fig. 14A and Fig. 14B show tumour growth and mouse weight for mice which received intra-tumoural injections, respectively. Tumour growth in mice which received carrier peptide or DMEM was greater than the untreated control mice group indicating the carrier peptide and culture medium provided a nutrient source for the tumours. In contrast, mice treated with the IK-1 peptide agent showed marked inhibition of tumour growth compared to the untreated control group. Substantially no difference in mouse weight was observed for the untreated and IK-1 treated mice groups at day 28 relative to time 0 (Fig. 14B). However, mice in the carrier peptide and DMEM treated groups showed marked weight loss compared to the control group at day 28. Inhibition of tumour growth was observed for mice injected intra-peritoneally with the peptide agent as shown in Fig. 15.

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

Dated: 17 October 2003

112266759

35.

Inter-K Pty Limited

Patent Attorneys for the Applicant:

BLAKE DAWSON WALDRON PATENT SERVICES

REFERENCES

- Agrez, M.V., Bates R.C., Mitchell, D., Wilson, N., Ferguson, N., Anseline, P. and Sheppard, D., Multiplicity of fibronectin-binding α v integrin receptors in colorectal cancer. *Br. J. Cancer* 73, 887-892 (1996).
- 5 Agrez, M.V., Gu, X., Turton, J., Meldrum, C., Niu, J., Antalis, T. and Howard, E.W., The α v β 6 integrin induces gelatinase B secretion in colon cancer cells. *Int. J. Cancer* 81, 90-97 (1999).
- Bachmann, A.S., Surovoy, A., Jung, G. and Moelling, K., Integrin receptor-targeted transfer peptides for efficient delivery of antisense oligodeoxynucleotides. *J. Mol. Med.* 76,
10 126-132 (1998).
- Boulton, T.G., Nye, S.H. and Robbins, D. J., ERKs: a family of protein-serine/ threonine kinases that are activated and tyrosine phosphorylated in response to insulin and NGF. *Cell* 65, 663-675 (1991).
- Breuss, J.M., Gallo, J., De Lisser, H.M., Klimanskaya, I.V., Folkesson, H.G., Pittet, J.F., Nishimura, S.L., Aldape, K., Landers, D.V., Carreter, W., Gillet, N., Sheppard, D.,
15 Mathay, M., Albeda, S.M., Kramer, R.H., and Pytela, R., Expression of the β 6 integrin in development, neoplasia and tissue repair suggests a role in epithelial remodelling. *J. Cell. Sci.* 108, 2241-2251 (1995).
- Breuss, J.M., Gillet, N., Lu, L., Sheppard, D. and Pytela, R., Restricted distribution
20 of integrin beta 6 mRNA in primate epithelial tissues. *J. Histochem. Cytochem.* 41, 1521-1527 (1993).
- Cone, R.I., Weinacker, A., Chen, A. and Sheppard, D., Effects of beta subunit cytoplasmic domain deletions on the recruitment of the integrin alpha v beta 6 to focal contacts. *Cell Adhes. Comm.* 2, 101-113 (1994).
- 25 Derossi, D., Calvet, S., Trembleau, A., et al, Cell internalization of the third helix of the Antennapedia homeo domain is receptor-independent. *J. Biol. Chem.* 271, 18188-18193 (1996).

- Derossi, D., Joliot, A.H., Chassaing, G. and Prochiantz, A., The third helix of the Antennapedia homeodomain translocates through biological membranes. *J. Biol. Chem.* 269, 10444-10450 (1994).
- 5 Garrington, T.P., and Johnson, G.L., Organization and regulation of mitogen-activated protein kinase signaling pathways. *Curr. Opin. Cell Biol.* 11, 211-218 (1999).
- Giancotti, F.G. and Ruoslahti, E., Integrin signalling. *Science* 285, 1028-1032 (1999).
- Haapasalmi, K., Zhang, K., Tonneson, M., Olerud, J., Sheppard, D., Salo, T., Kramer, R., Clark, R.A.F., Uitto, V.-J. and Larjava, H., Keratinocytes in human wounds express $\alpha 6 \beta 6$ integrin. *J. Invest. Dermatol.* 106, 42-48 (1996).
- 10 He, T.-C., Zhou, S., Da Costa, L.T., Yu, J., Kinzler, K.W. and Vogelstein, B. A simplified system for generating recombinant adenoviruses. *Proc., Natl. Acad. Sci. USA* 95, 2509-2514 (1998).
- Howe, A., Aplin, A.E., Alahari, S.K. and Juliano, R.L., Integrin signaling and cell growth control. *Curr. Opin. Cell Biol.* 10, 220-231 (1998)
- 15 Jones, J., Watt, F.M. and Speight, P.M., Changes in the expression of αv integrins in oral squamous cell carcinomas. *J. Oral Path. & Med.* 26, 63-68 (1997).
- Reszka, A.A., Hayashi, Y. and Horwitz, A.F., Identification of amino acid sequences in the integrin $\beta 1$ cytoplasmic domain implicated in cytoskeletal association. *J. Cell Biol.* 117, 1321-1330 (1992).
- 20 Schiller et al. *Int. J. Pept. Prot. Res.* 25, 175 (1985).
- Sheppard, D., Rozzo, C., Starr, L., Quaranta, V., Erle, D.J. and Pytela, R., Complete amino acid sequence of a novel integrin β subunit ($\beta 6$) identified in epithelial cells using the polymerase chain reaction. *J. Biol. Chem.* 265, 11502-11507 (1990).
- Smythe, W.R., Lebel, E., Bavaria, J.E., Kaiser, L.R. and Albelda, S.M., Integrin expression in non-small cell carcinoma of the lung. *Cancer & Metastasis Reviews* 14, 229-239 (1995).
- 25

Takiuchi, H., Kanokogi, M., Fujimoto, N., Hanafusa, T., Kyo, M., Ichikawa, Y., Nagano, S., Fukunishi, T., Yabumoto, H. and Ihara, H., Expression of integrin molecule in urological tumour cell lines by using RT-PCR method. Jap. J. Urology 85, 584-588 (1994).

- 5 Thomas, G.J., Jones, J. and Speight, P.M., Integrins and oral cancer. Oral Oncology 33, 381-388 (1997).

1/15




FIG. 1




FIG. 2

2/15




FIG. 3







FIG. 4

3/15



4/15

HUVEC Cell Line






FIG. 8A.


αV




$\beta 1$



$\beta 3$




$\beta 5$




$\beta 6$

SH-SY5Y cell line


FIG. 8B



$\beta 3$



$\beta 5$




$\beta 6$


5/15

HL 60 cell line


DU145 Cells



$\beta 3$



$\beta 5$



$\beta 6$

FIG. 8C








FIG. 8D

6/15


HMEC-1 CELL LINE



$\beta 3$



$\beta 5$



$\beta 6$

FIG. 8E

7/15

HT29 CELL LINE








FIG. 9A

H460 cell line



B5




B6


FIG. 9B

8/15


WM 115 cell line



$\beta 3$



$\beta 5$



$\beta 6$

FIG. 9C

9/15




FIG. 10A




FIG. 10B

10/15

Prostate Cancer (DU145)








FIG. 10D

11/15




Lung Cancer (H460)



Melanoma (WM115)



12/15



13/15





FIG. 13

14/15





FIG. 14B

15/15




FIG 15.

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/AU04/001416

International filing date: 15 October 2004 (15.10.2004)

Document type: Certified copy of priority document

Document details: Country/Office: AU
Number: 2003905726
Filing date: 17 October 2003 (17.10.2003)

Date of receipt at the International Bureau: 08 November 2004 (08.11.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.